



Climate change and occupational allergies: an overview on biological pollution, exposure and prevention

Maria Concetta D'Ovidio¹, Isabella Annesi-Maesano², Gennaro D'Amato³ and Lorenzo Cecchi⁴

¹Dipartimento di Medicina, Epidemiologia, Igiene del Lavoro e Ambientale, Istituto Nazionale per l'Assicurazione contro gli Infortuni sul Lavoro (INAIL), Monte Porzio Catone (Rome), Italy

²Sorbonne Universités, UPMC Univ Paris 06, INSERM, Pierre Louis Institute of Epidemiology and Public Health (IPLESP UMRS 1136), Epidemiology of Allergic and Respiratory Diseases Department (EPAR), Saint-Antoine Medical School, F75012 Paris, France

³Unità Operativa Complessa di Malattie Respiratorie e Allergiche, Azienda Ospedaliera ad Alta Specialità di Rilievo Nazionale A. Cardarelli, Naples, Italy

⁴Centro di Bioclimatologia, Università degli Studi di Firenze, Florence, Italy

Abstract

Introduction. Climate change, air pollution, temperature increase and other environmental variables are modifying air quality, contributing to the increase of prevalence of allergic respiratory diseases. Allergies are complex diseases characterized by multilevel interactions between individual susceptibility, response to immune modulation and environmental exposures to physical, chemical and biological agents. Occupational allergies introduce a further complexity to these relationships by adding occupational exposure to both the indoor and outdoor ones in the living environment.

Objective. The aim of this paper is to overview climate-related allergy affecting environmental and occupational health, as literature data are scanty in this regard, and to suggest a management model of this risk based on a multidisciplinary approach, taking the case of biological pollution, with details on exposure and prevention.

Conclusions. The management of climate-related occupational allergy should take into account preventive health strategies, environmental, public and occupational interventions, as well as to develop, implement, evaluate, and improve guidelines and standards protecting workers health under changing climatic conditions; new tools and strategies based on local conditions will have to be developed. Experimental studies and acquisition of environmental and personal data have to be matched to derive useful information for the scope of occupational health and safety.

Key words

- climate change
- allergy
- biological products
- worker
- air pollution

INTRODUCTION

Article 1 of the Framework Convention on Climate Change (UNFCCC) defines climate change as a *change of climate which is attributed directly or indirectly to human activity that alters the composition of the global atmosphere and which is in addition to natural climate variability observed over comparable time periods* [1].

Climate change in IPCC's (Intergovernmental Panel on Climate Change) position refers to a change in the state of the climate that can be identified (e.g. using statistical tests) by changes in the mean and/or the variability of its properties, and that persists for an extended period, typically decades or longer. It refers to any

change in climate over time, due to natural variability or as a result of human activity; it is responsible for increased temperature, rise in the sea level, modified air quality [2].

"Air pollution is the contamination of the indoor or outdoor environment by any chemical, physical or biological agent that modifies the natural characteristics of the atmosphere". Household combustion devices, motor vehicles, industrial facilities and forest fires are common sources of air pollution. Pollutants of major public health concern include particulate matter, carbon monoxide, ozone, nitrogen dioxide and sulfur dioxide. "Changes in temperature, modifications in pre-

precipitation frequency, and air stagnation also affect air pollution levels with significant risks to health. Climate affects pollution levels through pollutant formation, transport, dispersion, and deposition. For instance, fine particulate air pollution is estimated to be responsible for 7 million additional deaths globally in 2012, mainly due to respiratory and cardiovascular disease. Its effect is amplified by changes in ambient temperature, precipitation frequency, and air stagnation – all crucial for air pollutants formation, transport, dispersion and deposition”. Outdoor and indoor air pollution contributes to the increase of the prevalence of allergic respiratory diseases [3], exacerbating clinical features.

The 2015 Lancet Commission on Health and Climate Change “recommends that over the next 5 years, governments: Invest in climate change and public health research, monitoring, and surveillance to ensure a better understanding of the adaptation needs and the potential health co-benefits of climate mitigation at the local and national level” [4].

Occupational allergies are complex diseases because of the influence of environmental and climatic factors, the presence of allergens of different origin, the individual sensitization and immunological responses, the clinical manifestations: different and complementary medical skills are required for their management [5].

The topic *climate change* linked to the *occupational health* requires a particular attention by authorities in order to promote research, monitoring, surveillance, best practices, protective and preventive measures, starting with specific control program regarding occupational health and evaluation of environmental and climate factors. According to the scientific literature, thermal stress, injuries and fatalities are the main consequences of climate change in the workplaces [6, 7].

Allergic and respiratory diseases in relation to climate change have been extensively studied with regards to public health but little attention has been paid on climate-related allergic and respiratory diseases in the occupational sector, although some pertinent concerns are well known since several years [5, 8-14].

In Italy occupational health and safety are regulated by the Legislative Decree 81/08, implementing the EU directives on the improvement of health and safety at work. Article 28 (Title I - Common principles) of the Decree states that the employer has to assess all risks for health and safety of the workers. However, the allergic risk is explicitly mentioned in the Title X of the Decree (*Exposure to biological agents*) and particularly in the annex XLVI, where *possible allergic effects* are reported for 10 agents (2 parasites and 8 fungi) with respect to the 375 classified biological agents [15].

In this study we focused our attention on outdoor and indoor biological pollution, exposure and prevention related to climate change, with the aim to emphasize a topic – allergy – affecting environmental and occupational health and to remark the need to manage this risk in a multidisciplinary way. We also stress the importance of studying not only well known occupational allergens such as latex, house dust and flour, but also allergies caused by new indoor and outdoor sources of exposure.

PLANTS AND ANIMALS DERIVED AEROALLERGENS, CLIMATE CHANGE, ENVIRONMENTAL POLLUTION

Allergens and sensitizing agents are traditionally classified as high-molecular-weight (HMW) (glyco) proteins of animal and vegetable origin, mainly represented by grain dust (including mites), bakery dust, fish proteins, laboratory animals, bird proteins, natural rubber latex, enzymes (especially in detergents), mold proteins, vegetable gums, soy bean dust, cotton, coffee, and other seed dusts. Low-molecular weight (LMW) agents are metals, acid anhydrides, isocyanates, biocides, reactive dyes and others [16].

Although HMW proteins generally induce an IgE-response, involved in the pathogenesis of respiratory diseases (asthma, rhinitis, etc.), and LMW agents can causes T-cell dependent delayed-type reactions, the pathogenic mechanism of many of them is still unknown, for instance cases of occupational contact urticaria caused by parvalbumin, and respiratory diseases induced by some chemical agents, as reported in the literature [16-21].

The beginning and development of occupational allergic diseases are regulated by a complex interaction which includes endogenous and exogenous factors, individual susceptibility, exposure in living and working environments with different level, duration and frequency, etc. Moreover, the influence of climatic and environmental drivers on physical, chemical and biological pollutants also plays an important role in the onset of allergic diseases [8, 22-24].

A lot of allergens are affected by climatic and environmental variations, and many studies considered the effects of climate change and chemical pollution on the features and seasonality of pollen and plant flowering; influence on individual susceptibility, genetic variations, health consequences such as respiratory and allergic diseases have also been reported [9, 25, 26].

In this regard, a comprehensive and detailed biointeractive model to justify the increased prevalence of asthma and allergic diseases observed in recent decades [27-31] is still lacking.

Climate change and pollutants like greenhouse gases (GHGs), sulfur oxide (SO), particulate matter (PM), ozone (O₃), nitric oxide (NO) have pronounced negative effects on both environmental and human health. Fauna, plant species, pollen (including spatiotemporal dynamics of allergenic pollen) and other biological components are affected by climatic variation, atmospheric factors, meteorological conditions. All these factors determine an increased allergenic effects of both vegetable and animal components, inducing and/or exacerbating allergy and chronic respiratory diseases, as well as decreasing the lung function [31-42].

It is important to note that also indoor sources contribute to outdoor air pollution. Moreover, pollution arising from chemical, physical and biological agents affects climatic parameters (temperature, humidity, wind speed etc.), but the opposite is also true in both workplaces and living environments.

Institutional networks and associations ensure a continuous monitoring of pollen, fungal species distri-

Table 1

An overview of some institutional links regarding climate change, air pollution, public health and occupational health

Institution	Link
Agenzia Regionale per la Protezione Ambientale (ARPA)	web-sites
Associazione Italiana di Aerobiologia (AIA)	www.ilpolline.it
Centro Euro Mediterraneo sui Cambiamenti Climatici (CMCC)	www.cmcc.it
Centers for Disease Control and Prevention (CDC)	www.cdc.gov
Centro Interdipartimentale di Bioclimatologia (CIBIC)	www.cibic.unifi.it
Centro nazionale per la prevenzione e il Controllo delle Malattie (CCM)	www.ccm-network.it
Consiglio Nazionale delle Ricerche (CNR)	www.cnr.it
Ente Nazionale per l'Energia e l'Ambiente (ENEA)	www.enea.it
European Center for Disease Prevention and Control (ECDC)	http://ecdc.europa.eu
European Environment Agency (EEA)	www.eea.europa.eu
Global Framework for Climate Services (GFCS)	http://gfcs-climate.org
Intergovernmental Panel on Climate Change (IPCC)	www.ipcc.ch
International Labour Organization (ILO)	www.ilo.org
Istituto Nazionale per l'Assicurazione contro gli Infortuni sul Lavoro (INAIL)	www.inail.it
Istituto Superiore di Sanità (ISS)	www.iss.it
Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA)	www.isprambiente.gov.it
Ministero dell'Ambiente e della Tutela del Territorio e del Mare	www.minambiente.it
Ministero della Salute	www.salute.gov.it
National Institute for Occupational Safety and Health (NIOSH)	www.osha.gov
Occupational Exposure to Hazards Agents (HazMap)	http://hazmap.nlm.nih.gov
Occupational Safety and Health Administration (OSHA)	www.osha.gov
United Nations Framework Convention Climate Change (UNFCCC)	www.unfccc.int
World Allergy Organization (WAO)	www.worldallergy.org
World Health Organization (WHO)	www.who.int
World Meteorological Organization (WMO)	www.wmo.int

bution and other biological emissions, as well as of parameters regarding climate and the environment (Table 1). In this perspective, the integration of environmental and climatic data as well as the synergy between different institutions in order to promote the protection of public and occupational health should be highly beneficial.

IMPACT OF CLIMATE CHANGE ON OCCUPATIONAL EXPOSURE

Outdoor workers are at high risk for exposure to allergens of biological (animal and vegetable) origin, because of the prolonged permanence in the workplace or the intense physical activity characterizing some working tasks resulting in an increased respiratory inhalation. Indoor environment can also be affected by climatic variations, as some environmental parameters and various types of pollutants increases the interchange, making recognition of a clear separation between indoor and outdoor difficult.

Several criticisms are inherent to occupational allergies of biological origin, such as the identification of all the biological agents with potential allergenic effects, since current regulation (in Italy the Legislative Decree 81/08 - annex XLVI: *Biological agents classified*) recog-

nizes only ten biological agents with regard to allergic effects (Table 2); moreover, evaluation of the role of several co-factors involved in both the individual susceptibility and allergic responses is still lacking. In addition several professional categories are potentially exposed to allergic risk. Construction, agriculture, animal husbandry, fishing, hunting, park maintenance, farming, forestry, urban rescue and safety are representative of outdoor workplaces; therefore subjects operating in the urban (ecological operators, safety and rescue operators, sport instructors, gardeners etc.) and suburban (farmers, ranchers, gardeners, beekeepers, foresters, safety's operators, surveillance activity etc.) environments can be defined outdoor workers. In this context risk assessment shall be performed, taking into account environmental data arising from physical, chemical and biological parameters as well as from results of health surveillance of exposed workers.

Occupational allergies have been traditionally studied in indoor environment (*i.e.* offices, schools, bakeries, hospitals) where main allergens are mites, dusts, latex, and other (micro)components [43-46]. In most cases, different methodologies have been applied to evaluate both the environmental exposure (air and sampling monitoring) and the immune response (skin tests,

Table 2

List of the classified biological agents with possible allergic effects

Classified biological agents = 375*		
Bacteria		N = 151
Viruses		N = 129
Parasites		N = 69
Fungi		N = 26
Biological agents with possible allergic effects		N = 10
Parasites		
Biological agents	Classification	Notes**
<i>Ascaris lumbricoides</i>	2	A
<i>Ascaris suum</i>	2	A
Fungi		
Biological agents	Classification	Notes**
<i>Aspergillus fumigatus</i>	2	A
<i>Candida albicans</i>	2	A
<i>Coccidioides immitis</i>	3	A
<i>Cryptococcus neoformans</i> var. <i>neoformans</i> (<i>Filobasidiella neoformans</i> var. <i>neoformans</i>)	2	A
<i>Cryptococcus neoformans</i> var. <i>gattii</i> (<i>Filobasidiella bacillispora</i>)	2	A
<i>Epidermophyton floccosum</i>	2	A
<i>Microsporum</i> spp.	2	A
<i>Penicillium marneffei</i>	2	A

*According to Annex XLVI – Legislative Decree 81/08

**A: Possible allergic effects

immunochemical methods, molecular assays) [47-49].

In order to contribute to a better understanding of occupational allergies, a bibliometric approach for the study of scientific literature was followed. The search criteria are shown in the Supplementary data 1-8 available online (*Supplementary Materials*); we used MeSH (*Medical Subject Headings*) Terms – the NLM controlled vocabulary thesaurus used for indexing articles for PubMed – regarding air pollutants or allergens or pollen or changes, climate, occupational health or occupational exposure, hypersensitivity or allergy and immunology or respiratory tract diseases. A limit of data searching was the lack of “outdoor” term. Meva (*MEDLINE Evaluator*) – a free medico-scientific data mining web service to analyse bibliographic MEDLINE data – was used as post-processor condensing the endless list of a MEDLINE result by presenting counts, frequency distributions, contingency tables (relations) and detailed sorted lists of selected bibliographic fields. We applied this bibliometric approach to the literature of the last 5 years. As we reported in a previous communication, the study of bibliometric sources showed that articles refer specifically to a particular allergic condition, a source of exposure to specific allergens (both vegetables and animals), preventive measures against specific occupational allergies. It's easy to extrapolate articles that deal with occupational allergies about indoor because this term is present in MeSH Terms, while terms outdoor/external/outer do not appear [50].

Workplaces may be an important source of exposure

to air pollution, both outdoor and indoor, not only for dust and fumes, but also for vegetable and animal allergens; examples include arthropod allergens such as house dusts, mites and cockroaches, mammalian allergens derived from pets or pests, mouse, rats, horses, jellyfish, hymenoptera, wild and domestic species such as wasps, bees, hornets, etc. [43-46, 51-58].

Workers or working activities at risk of exposure to biological allergens are represented by flour millers, pastry makers, health-care workers, laboratory technicians, farmers, sea foods processing, baking products manufacturing detergent production, pharmaceutical industry, food industry, foresters, beekeepers, marine biologists, fishermen, scuba divers, gardeners, farmers, firefighters, foresters, municipal operators. Following exposure to allergen, clinical features may involve respiratory tract (occupational asthma is the most frequent work-related disease), eye and skin tissues, other systems following ingestion, contact and inoculation of the allergens. The management needs a multidisciplinary approach through innovative molecular methodologies applied to both environmental and biological matrices, medical history acquisition, health surveillance and control measures regarding outdoor and indoor environmental exposures [59-63].

The management of occupational allergy should take into account all factors regarding lifestyles, including hobbies and outdoor recreational activities, as well as indirect, no apparent exposure or passive carriage of

allergens [64-66]. It is also necessary to assess the co-exposure to other environmental factors such as solar radiation [67] that can modulate the immune response and the occurrence of symptoms at various grades of severity. Moreover it should be noted that *"one health frames the complex interactions between human, animal and environmental health, one health is a unique approach to cope with allergies, one health focuses on equity and is key for human sustainable development"* [68].

COLLABORATIVE, MULTIDISCIPLINARY AND INTEGRATED APPROACH

In public and occupational settings health prevention and promotion are intrinsically linked to education through information, training and communication. It is important to increase the awareness of workers regarding the risk derived from exposure to allergens and to define collective and individual control measures. Moreover, workers may be responsible for the spread of allergens to the community, carrying allergens through their clothes, hair and shoes. The innovative molecular methodologies on biological fluids that allow the determination of many allergens with a high degree of sensitivity are very useful if implemented during workers' health surveillance. It is likely that the development of molecular aerobiology will provide a more detailed approach in term of health impacts. A multidisciplinary approach is highly advisable to study the full spectrum of issues regarding climate change, aerobiology and clinical outcomes [69, 70].

Preventive health strategies should involve environmental, public and occupational interventions. Decision makers have the responsibility to develop, implement, evaluate, and improve guidelines and standards to protect workers' health under changing climatic conditions; new tools and strategies based on local conditions will have to be developed.

As shown in Figure 1, Figure 2a and Figure 2b (available online as *Supplementary Materials*) the study of biological occupational allergies should be performed by a multidisciplinary approach based on both traditional and innovative tools. Regarding exposure to vegetable allergens – pollens in particular – air monitoring is addressed to improve the new molecular approaches in addition to traditional methodology used by monitoring stations [69,70].

Starting from a list of allergens [20] it is necessary to know all information to manage the risk due to occupational allergy, such as sampling methodologies that should be standardized. In order to protect the health of both workers and the general public and to safeguard plants and animals, the development of new and innovative methods for sampling and measurement of (bio) pollutants should be a priority.

Our research is addressed to carry out strategies and programs for the management of outdoor occupational allergy, listed in following key points (Figure 1, Figure 2a and Figure 2b, available online as *Supplementary Materials*):

- I. Climate change and occupational allergy diseases (CCOADs)
- II. Allergy program control (APC)

III. Personal protective equipment (PPE)

IV. Communication, education and training (CET)

V. General and scientific information (GSI)

VI. Occupational and non occupational fields (ONOFs)

VII. Forecasting Models (FoMs)

VIII. Information and communication technology (ICT)

Consultation of the national and international literature supported by a bibliometric analysis confirmed the lack of research on the assessment of biological allergens in outdoor workplaces. This analysis provides information for research teams and audience to clearly identify or mark peculiar topics in investigated fields.

For the protection of public health, the presence of monitoring networks allows the sampling of pollen and fungal spores through techniques fundamentally based on the principles of gravitational deposition, inertial impact and sucking, using stationary samplers placed at certain heights above ground that allow sampling on an ongoing basis. The measurement of outdoor allergens is based on standardized methods (reading on a microscope slide) and new molecular technologies are currently being tested. The standardization of sampling aerobiological methodology should also be conducted at national and international level. As part of aerobiology some goals to be pursued include the harmonization of existing networks for monitoring biological particles, the development of strategies to combat pollution and the identification of the role of climate change in allergic diseases for public and occupational health purposes. It is necessary to define the role of the environment in the prevention of allergic diseases, the factors affecting levels of indoor and outdoor allergens and the methods to reduce exposure in the living and working environments, taking into account both the individual and environment exposure [5, 22, 23, 69-76].

Research topics should include action mechanisms, classification, prevention, epidemiological surveillance/registries, predictive biomarkers, molecular diagnosis, treatment response, phenotypes, serious illness/exacerbations, new treatments (biological vaccine and drugs). Data collected through the environmental networks should be matched with those concerning workers health. *Ad hoc* questionnaires, immunochemical and molecular tests to assess immunological and other physiological parameters are of paramount importance to plan appropriate technical, organizational and procedural interventions addressed at specific workplaces. Information, education and communication are unreplaceable tools.

Our research aim is to perform strategies and programs for the management of outdoor and indoor occupational allergy through an experimental, integrated and multidisciplinary approach, also considering mathematical models as valid tools to analyze emission and dispersion phenomena, contributing to design risk scenarios.

CONCLUDING REMARKS

Allergy is a complex disease characterized by multiple interactions between individual susceptibility, genetic background, response to immune modulation and environmental exposures to physical, chemical and biological agents. Occupational allergies amplify these

relationships by adding other factors, depending on the workplace exposures. Traditionally, the study of occupational allergies was focused on indoor pollutants, but the growing body of data about allergies in the general population and recent new evidences are a stimulus to extend research to outdoor environment, trying to understand the potential connection between indoor and outdoor allergens. Air pollution and climate change have a central role in the etiology and pathology of occupational allergies.

A great body of literature suggests that bibliometric indicators are one of the most common tools to measure research performance. Moreover, these indicators have a wide range of applications, such as *descriptive linguistics*, development of *thesauri*, evaluation of reader usage and measuring term frequencies.

It is necessary to promote a multidisciplinary approach starting from an exhaustive and detailed knowledge of the literature regarding allergies, together with the monitoring of physical, chemical and biological parameters, data on the immune-susceptibility of the workers and the influence of other co-factors. In addition, forecasting and mathematical models can be helpful to increase the knowledge and to suggest the preventive measures to adopt in this area.

In conclusion, climate change represents an issue that need to be explored and studied in its different and complex aspects. The impact resulting from temperature increase and air pollution, health consequences such as cardiovascular and respiratory diseases and accidents and other events is certainly recognized.

These features are the apical part of a phenomenon

which – if not counteracted – can keep on and greatly damage the environment and human health. Local actions as well as national and international agreements are aimed at safeguarding the “universal health”, since environmental and human health are interconnected. The questions regarding occupational health should be considered in an integrated life course approach [76-78].

Acknowledgements

We are grateful to Nunzia Bellantonio, Raffaella Modestino and Stefania Massari for their helpful collaboration and discussion regarding bibliometric approach, Claudio Gariazzo for his helpful collaboration and discussion concerning forecasting models, and Nicoletta Vonesch for the general revision. Availability, courtesy and professionalism are qualities that we wish to highlight.

This paper is part of a monographic section dedicated to Climate change and occupational health, edited by Maria Concetta D'Ovidio, Carlo Grandi, Enrico Marchetti, Alessandro Polichetti and Sergio Iavicoli and published in the same issue: *Ann Ist Super Sanità* 2016;52(3):323-423.

Conflict of interest statement

There are no potential conflicts of interest or any financial or personal relationships with other people or organizations that could inappropriately bias conduct and findings of this study.

Submitted on invitation.

Accepted on 12 April 2016.

REFERENCES

1. United Nations. *United Nations Framework Convention on Climate Change* (UNFCCC). New York City, US, 9 May, 1992. Available from: <http://unfccc.int/2860.php>.
2. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. *Climate Change 2014. Impacts, adaptation, and vulnerability. Part A. Global and sectoral aspects*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2014.
3. World Health Organization. *Burden on disease from air pollution in 2012*. Geneva: WHO; 2014.
4. Watts N, Adger WN, Agnolucci P, Blackstock J, Byass P, Cai W, Chaytor S, Colbourn T, Collins M, Cooper A, Cox PM, Depledge J, Drummond P, Ekins P, Galaz V, Grace D, Graham H, Grubb M, Haines A, Hamilton I, Hunter A, Jiang X, Li M, Kelman I, Liang L, Lott M, Lowe R, Luo Y, Mace G, Maslin M, Nilsson M, Oreszczyn T, Pye S, Quinn T, Svensdotter M, Venevsky S, Warner K, Xu B, Yang J, Yin Y, Yu C, Zhang Q, Gong P, Montgomery H, Costello A. Health and climate change: policy responses to protect public health. *Lancet* 2015;386(10006):1861-914. DOI: 10.1016/S0140-6736(15)60854-6
5. Raulf M, Buters J, Chapman M, Cecchi L, de Blay F, Doekes G, Eduard W, Heederik D, Jeebhay MF, Keespohl S, Krop E, Moscato G, Pala G, Quirce S, Sander I, Schlünssen V, Sigsgaard T, Walusiak-Skorupa J, Wisniewska M, Wouters IM, Annesi-Maesano I. European Academy of Allergy and Clinical Immunology. Monitoring of occupational and environmental aeroallergens-EAACI Position Paper. Concerted action of the EAACI IG Occupational Allergy and Aerobiology & Air Pollution. *Allergy* 2014;69(10):1280-99. DOI: 10.1111/all.12456
6. Oyekale AS. Climate change induced occupational stress and reported morbidity among cocoa farmers in South-Western Nigeria. *Ann Agric Environ Med* 2015;22(2):357-61. DOI: 10.5604/12321966.1152095
7. Tawatsupa B, Yiengprugsawan V, Kjellstrom T, Berecki-Gisolf J, Seubsman SA, Sleight A. Association between heat stress and occupational injury among Thai workers: findings of the Thai Cohort Study. *Ind Health* 2013;51(1):34-46.
8. D'Amato G, Holgate ST, Pawankar R, Ledford DK, Cecchi L, Al-Ahmad M, Al-Enezi F, Al-Muhsen S, Ansotegui I, Baena-Cagnani CE, Baker DJ, Bayram H, Bergmann KC, Boulet LP, Buters JT, D'Amato M, Dorsano S, Douwes J, Finlay SE, Garrasi D, Gómez M, Haahtela T, Halwani R, Hassani Y, Mahboub B, Marks G, Michelozzi P, Montagni M, Nunes C, Oh JJ, Popov TA, Portnoy J, Ridolo E, Rosário N, Rottem M, Sánchez-Borges M, Sibanda E, Sienra-Monge JJ, Vitale C, Annesi-Maesano I. Meteorological conditions, climate change, new emerging factors, and asthma and related allergic disorders. A statement of the World Allergy Organization. *World Allergy Organ J* 2015;8(1):25. eCollection 2015. DOI:

- 10.1186/s40413-015-0073-0
9. D'Amato G, Bergmann KC, Cecchi L, Annesi-Maesano I, Sanduzzi A, Liccardi G, Vitale C, Stanzola A, D'Amato M. Climate change and air pollution. Effects on pollen allergy and other allergic respiratory diseases. *Allergo J Int* 2014;23:17-23. DOI 10.1007/s40629-014-0003-7
10. Cecchi L, D'Amato G, Annesi Maesano I. Climate, urban air pollution, and respiratory allergy. *Climate vulnerability: understanding and addressing threats to essential resources*. Elsevier Inc., Academic Press, 105-113, 2013.
11. D'Amato G, Baena-Cagnani CE, Cecchi L, Annesi-Maesano I, Nunes C, Ansotegui I, D'Amato M, Liccardi G, Sofia M, Canonica WG. Climate change, air pollution and extreme events leading to increasing prevalence of allergic respiratory diseases. *Multidiscip Respir Med* 2013; 8(1):12. DOI: 10.1186/2049-6958-8-12
12. D'Ovidio MC, Grandi C. Cambiamento climatico, inquinamento atmosferico, allergie occupazionali aero-trasmesse. In: 78° Congresso Nazionale SIMLII (Società Italiana di Medicina del Lavoro ed Igiene Industriale), 25-27 Novembre 2015, Milano. *Med Lav* 2015;106(Suppl 2):81.
13. D'Ovidio MC, Tomao P, Vonesch N. Cambiamenti climatici, disastri naturali: maggior rischio di esposizione ad agenti biologici e allergologici per gli operatori dell'emergenza. In: XV Giornata mondiale dell'acqua - Convegno "Grado di inquinamento naturale di acque e suoli in Italia". *Accademia Nazionale dei Lincei*. Roma 20 marzo 2015.
14. Sidi E, Albahary C, Longueville R. Allergy and occupational diseases. *Ind Med Surg* 1953;22(3):93-8.
15. Italia. Decreto Legislativo 9 aprile, n. 81. Attuazione dell'articolo 1 della legge 3 agosto 2007, n. 123, in materia di tutela della salute e della sicurezza nei luoghi di lavoro. *Gazzetta Ufficiale* n. 101, 30 aprile 2008. *Suppl. Ord.* n. 108L.
16. Vandenplas O, Worm M, Cullinan P, Park HS, van Wijk RG. Section 2.10. Occupational Allergy. In: Pawankar R, Canonica GW, Holgate ST, Lockey RF, Blaiss MS (Eds). *White Book on Allergy: Update 2013*. *World Allergy Organization*; 2013. p. 72-77.
17. Sano A, Yagami A, Suzuki K, Iwata Y, Kobayashi T, Arima M, Kondo Y, Yoshikawa T, Matsunaga K. Two cases of occupational contact urticaria caused by percutaneous sensitization to parvalbumin. *Case Rep Dermatol* 2015;7(2):227-32. DOI: 10.1159/000439080. Erratum in: *Case Rep Dermatol* 2015;7(3):328.
18. Seed MJ, Enoch SJ, Agius RM. Chemical determinants of occupational hypersensitivity pneumonitis. *Occup Med* 2015;65(8):673-81. DOI: 10.1093/occmed/kqv143
19. Laštovková A, Klusáčková P, Fenclová Z, Bonnetterre V, Pelclová D. Asthma caused by potassium aluminium tetrafluoride: a case series. *Ind Health* 2015;53(6):562-8. DOI: 10.2486/indhealth.2014-0094
20. Baur X, Bakehe P. Allergens causing occupational asthma: an evidence-based evaluation of the literature. *Int Arch Occup Environ Health* 2014;87(4):339-63. DOI: 10.1007/s00420-013-0866-9
21. Barbaud A, Poreaux C, Penven E, Waton J. Occupational protein contact dermatitis. *Eur J Dermatol* 2015;25(6):527-34. DOI: 10.1684/ejd.2015.2593
22. Peden D, Reed CE. Environmental and occupational allergies. *J Allergy Clin Immunol* 2010;125(2 Suppl 2):S150-160. DOI: 10.1016/j.jaci.2009.10.073
23. Zissler UM, Esser-von Bieren J, Jakwerth CA, Chaker AM, Schmidt-Weber CB. Current and future biomarkers in allergic asthma. *Allergy* 2016;71(4):475-94. DOI: 10.1111/all.12828
24. Beasley R, Semprini A, Mitchell EA. Risk factors for asthma: is prevention possible? *Lancet* 2015;386(9998):1075-85. DOI: 10.1016/S0140-6736(15)00156-7
25. Abrahamczyk S, Kluge J, Gareca Y, Reichle S, Kessler M. The influence of climatic seasonality on the diversity of different tropical pollinator groups. *PLoS One* 2011;6(11):e27115. DOI: 10.1371/journal.pone.0027115
26. D'Amato G, Vitale C, De Martino A, Viegi G, Lanza M, Molino A, Sanduzzi A, Vatrella A, Annesi-Maesano I, D'Amato M. Effects on asthma and respiratory allergy of climate change and air pollution. *Multidiscip Respir Med* 2015;10:39. DOI: 10.1186/s40248-015-0036-x
27. Platts-Mills TA. The allergy epidemics: 1870-2010. *J Allergy Clin Immunol* 2015;136(1):3-13. DOI: 10.1016/j.jaci.2015.03.048
28. Yoo B, Park Y, Park K, Kim H. A 9-year trend in the prevalence of allergic disease based on national health insurance data. *J Prev Med Public Health* 2015;48(6):301-9. DOI: 10.3961/jpmph.15.011
29. An SY, Choi HG, Kim SW, Park B, Lee JS, Jang JH, Sung MW. Analysis of various risk factors predisposing subjects to allergic rhinitis. *Asian Pac J Allergy Immunol* 2015;33(2):143-51. DOI: 10.12932/AP0554.33.2.2015
30. Huang SK, Zhang Q, Qiu Z, Chung KF. Mechanistic impact of outdoor air pollution on asthma and allergic diseases. *J Thorac Dis* 2015;7(1):23-33. DOI: 10.3978/j.issn.2072-1439.2014.12.13 Erratum in: *J Thorac Dis* 2015;7(10):E521.
31. Baldacci S, Maio S, Cerrai S, Sarno G, Baiz N, Simoni M, Annesi-Maesano I, Viegi G; HEALS Study. Allergy and asthma: effects of the exposure to particulate matter and biological allergens. *Respir Med* 2015;109(9):1089-104. DOI: 10.1016/j.rmed.2015.05.017
32. Zhang Y, Bielory L, Mi Z, Cai T, Robock A, Georgopoulos P. Allergenic pollen season variations in the past two decades under changing climate in the United States. *Glob Chang Biol* 2015;21(4):1581-9. DOI: 10.1111/gcb.12755
33. Beck I, Jochner S, Gilles S, McIntyre M, Buters JT, Schmidt-Weber C, Behrendt H, Ring J, Menzel A, Traidl-Hoffmann C. High environmental ozone levels lead to enhanced allergenicity of birch pollen. *PLoS One* 2013;8(11):e80147. DOI: 10.1371/journal.pone.0080147
34. D'Amato G, Rottem M, Dahl R, Blaiss M, Ridolo E, Cecchi L, Rosario N, Motala C, Ansotegui I, Annesi-Maesano I, WAO Special Committee on Climate Change and Allergy. Climate change, migration, and allergic respiratory diseases: an update for the allergist. *World Allergy Organ J* 2011;4(7):120-5.
35. Jalbert I, Golebiowski B. Environmental aeroallergens and allergic rhino-conjunctivitis. *Curr Opin Allergy Clin Immunol* 2015;15(5):476-81. DOI: 10.1097/ACI.0000000000000205
36. Damialis A, Mohammad AB, Halley JM, Gange AC. Fungi in a changing world: growth rates will be elevated, but spore production may decrease in future climates. *Int J Biometeorol* 2015;59(9):1157-67. DOI: 10.1007/s00484-014-0927-0
37. Tankersley MS, Ledford DK. Stinging insect allergy: state of the art 2015. *J Allergy Clin Immunol Pract* 2015;3(3):315-22. DOI: 10.1016/j.jaip.2015.03.012
38. Butt N, Seabrook L, Maron M, Law BS, Dawson TP, Syktus J, McAlpine CA. Cascading effects of climate extremes on vertebrate fauna through changes to low-latitude tree flowering and fruiting phenology. *Glob Chang Biol* 2015;21(9):3267-77. DOI: 10.1111/gcb.12869
39. Favero-Longo SE, Sandrone S, Matteucci E, Appolonia L, Piervittori R. Spores of lichen-forming fungi in the myco-aerosol and their relationships with climate factors. *Sci Total Environ* 2014;466-467:26-33. DOI: 10.1016/j.scito-

- tenv.2013.06.057
40. Le Conte Y, Navajas M. Climate change: impact on honey bee populations and diseases. *Rev Sci Tech* 2008;27(2):485-97, 499-510.
 41. Zhang Q, Qiu Z, Chung KF, Huang SK. Link between environmental air pollution and allergic asthma: East meets West. *J Thorac Dis* 2015;7(1):14-22. Erratum in: *J Thorac Dis* 2015;7(10):E520. DOI: 10.3978/j.issn.2072-1439.2014.12.07
 42. Lang-Yona N, Levin Y, Dannemiller KC, Yarden O, Pecia J, Rudich Y. Changes in atmospheric CO₂ influence the allergenicity of *Aspergillus fumigatus*. *Glob Chang Biol* 2013;19(8):2381-8. DOI: 10.1111/gcb.12219
 43. Raulf M, Bergmann KC, Kull S, Sander I, Hilger C, Brüning T, Jappe U, Müsken H, Sperl A, Vrtala S, Zahradnik E, Klimek L. Mites and other indoor allergens – from exposure to sensitization and treatment. *Allergo J Int* 2015;24(3):68-80.
 44. Liss GM, Sussman GL, Deal K, Brown S, Cividino M, Siu S, Beezhold DH, Smith G, Swanson MC, Yunginger J, Douglas A, Holness DL, Lebert P, Keith P, Wasserman S, Turjanmaa K. Latex allergy: epidemiological study of 1351 hospital workers. *Occup Environ Med* 1997;54(5):335-42.
 45. Norbäck D, Michel I, Widström J. Indoor air quality and personal factors related to the sick building syndrome. *Scand J Work Environ Health* 1990;16(2):121-8.
 46. Gravesen S, Larsen L, Gyntelberg F, Skov P. Demonstration of microorganisms and dust in schools and offices. An observational study of non-industrial buildings. *Allergy* 1986;41(7):520-5.
 47. Lim FL, Hashim Z, Than LT, Md Said S, Hisham Hashim J, Norbäck D. Asthma, airway symptoms and rhinitis in office workers in Malaysia: associations with house dust mite (HDM) allergy, cat allergy and levels of house dust mite allergens in office dust. *PLoS One* 2015;10(4):e0124905. DOI: 10.1371/journal.pone.0124905
 48. Pesonen M, Jolanki R, Laresse Filon F, Wilkinson M, Kręcis B, Kieć-Swierczyńska M, Bauer A, Mahler V, John SM, Schnuch A, Uter W; ESSCA network. Patch test results of the European baseline series among patients with occupational contact dermatitis across Europe - analyses of the European Surveillance System on Contact Allergy network, 2002-2010. *Contact Dermatitis* 2015;73(3):163-71. DOI: 10.1111/cod.12333
 49. Jang JH, Kim DW, Kim SW, Kim DY, Seong WK, Son TJ, Rhee CS. Allergic rhinitis in laboratory animal workers and its risk factors. *Ann Allergy Asthma Immunol* 2009;102(5):373-7. DOI: 10.1016/S1081-1206(10)60507-8
 50. Bellantonio N, Modestino R, Wirz A, Riviello MC, Masari S, D'Ovidio MC. Revisione della letteratura riguardante gli effetti sulla salute occupazionale causati dalla dispersione di allergeni di origine vegetale e animale. In: XXXIX Congresso AIE, Alimentazione e salute nell'era della globalizzazione. Milano 28-30 ottobre 2015. Poster - 031.
 51. Karvala K, Toskala E, Luukkonen R, Uitti J, Lappalainen S, Nordman H. Prolonged exposure to damp and moldy workplaces and new-onset asthma. *Int Arch Occup Environ Health* 2011;84(7):713-21. DOI: 10.1007/s00420-011-0677-9
 52. Bush RK, Stave GM. Laboratory animal allergy: an update. *ILAR J* 2003;44(1):28-51.
 53. D'Ovidio MC, Martini A, Melis P, Signorini S. Value of the microarray for the study of Laboratory Animal Allergy (LAA). *G Ital Med Lav Ergon* 2011;33(2):109-16.
 54. Oldenburg M, Petersen A, Baur X. Maize pollen is an important allergen in occupationally exposed workers. *J Occup Med Toxicol* 2011;6:32. DOI: 10.1186/1745-6673-6-32
 55. Moscato G, Pala G, Crivellaro M, Siracusa A. Anaphylaxis as occupational risk. *Curr Opin Allergy Clin Immunol* 2014;14(4):328-33. DOI: 10.1097/ACI.0000000000000066
 56. Suarathana E, Shen A, Henneberger PK, Kreiss K, Leppla NC, Bueller D, Lewis DM, Bledsoe TA, Janotka E, Peterson EL. Post-hire asthma among insect-rearing workers. *J Occup Environ Med* 2012;54(3):310-7. DOI: 10.1097/JOM.0b013e31823fe098
 57. Majasuo S, Liippo J, Lammintausta K. Non-occupational contact sensitization to epoxy resin of bisphenol A among general dermatology patients. *Contact Dermatitis* 2012;66(3):148-53. DOI: 10.1111/j.1600-0536.2011.01993.x
 58. Liccardi G, Salzillo A, Dente B, Piccolo A, Lobefalo G, Noschese P, Russo M, Gilder JA, D'Amato G. Horse allergens: an underestimated risk for allergic sensitization in an urban atopic population without occupational exposure. *Respir Med* 2009;103(3):414-20. DOI: 10.1016/j.rmed.2008.09.017
 59. Moscato G, Maestrelli P, Bonifazi F, Troise C, Caminati M, Crivellaro M, Olivieri M, Senna G. OCCUPATION study (OCCUPation asthma: a nAtional based study): a survey on occupational asthma awareness among Italian allergists. *Eur Ann Allergy Clin Immunol* 2014;46(1):26-9.
 60. Tan J, Bernstein JA. Occupational asthma: an overview. *Curr Allergy Asthma Rep* 2014;14(5):431. DOI: 10.1007/s11882-014-0431-y
 61. Tarlo S. Clinical aspects of work-related asthma: past achievements, persistent challenges, and emerging triggers. *J Occup Environ Med* 2014;56(Suppl 10):S40-4. DOI: 10.1097/JOM.0000000000000285
 62. Boonchai W, Sirikudta W, Iamtharachai P, Kasemsarn P. Latex glove-related symptoms among health care workers: a self-report questionnaire-based survey. *Dermatitis* 2014;25(3):135-9. DOI: 10.1097/DER.0000000000000044
 63. Rask-Andersen A. Asthma increase among farmers: a 12-year follow-up. *Ups J Med Sci* 2011;116(1):60-71. DOI: 10.3109/03009734.2010.503287
 64. Liccardi G, Baldi G, Ciccarello A, Cutajar M, D'Amato M, Gargano D, Giannattasio D, Leone G, Schiavo ML, Madonna F, Menna G, Montera C, Pio A, Russo M, Salzillo A, Stanziola A, D'Amato G. Sensitization to rodents (mouse/rat) in urban atopic populations without occupational exposure living in Campania district (Southern Italy): a multicenter study. *Multidiscip Respir Med* 2013;8(1):30. DOI: 10.1186/2049-6958-8-30
 65. Liccardi G, Piccolo A, Dente B, Salzillo A, Noschese P, Gilder JA, Russo M, D'Amato G. Rabbit allergens: a significant risk for allergic sensitization in subjects without occupational exposure. *Respir Med* 2007;101(2):333-9.
 66. Liccardi G, Dente B, Senna G, De Martino M, D'Amato L, D'Amato G. Sensitization to horse allergens without apparent exposure to horse. Report of two cases. *Eur Ann Allergy Clin Immunol* 2005;37(9):350-2.
 67. D'Ovidio MC, Grandi C. Solar radiation and the immune system: concerns for outdoor workers. In: *Proceedings of ICOH 31st International Congress on Occupational Health*. May 31-June 5, 2015 Seoul, Korea. Abstract n. IC-0059.
 68. Cezmi AA, Agache I (Eds). *Global atlas of allergy*. European Academy of Allergy and Clinical Immunology (EAACI); 2014.
 69. Venanzetti F, D'Ovidio MC. Allergie di origine vegetale e qualità dell'aria: utilità delle tecniche molecolari per l'identificazione tassonomica di pollini per la tutela della salute negli ambienti di lavoro. In: XXXIX Congresso AIE, Alimentazione e salute nell'era della globalizzazione. Milano, 28-30 ottobre 2015. Poster - 032.

70. Cecchi L. From pollen count to pollen potency: the molecular era of aerobiology. *Eur Respir J* 2013;42(4):898-900. DOI: 10.1183/09031936.00096413
71. Amaral AF, Newson RB, Abramson MJ, Antó JM, Bono R, Corsico AG, de Marco R, Demoly P, Forsberg B, Gislason T, Heinrich J, Huerta I, Janson C, Jögi R, Kim JL, Maldonado J, Martinez-Moratalla Rovira J, Neukirch C, Nowak D, Pin I, Probst-Hensch N, Raheison-Semjen C, Svanes C, Urrutia Landa I, van Ree R, Versteeg SA, Weyler J, Zock JP, Burney PG, Jarvis DL. Changes in IgE sensitization and total IgE levels over 20 years of follow-up. *J Allergy Clin Immunol* 2016;137(6):1788-1795.e9. DOI: 10.1016/j.jaci.2015.09.037
72. Brauer M, Freedman G, Frostd J, van Donkelaar A, Martin RV, Dentener F, Dingenen Rv, Estep K, Amini H, Apte JS, Balakrishnan K, Barregard L, Broday D, Feigin V, Ghosh S, Hopke PK, Knibbs LD, Kokubo Y, Liu Y, Ma S, Morawska L, Sangrador JL, Shaddick G, Anderson HR, Vos T, Forouzanfar MH, Burnett RT, Cohen A. Ambient air pollution exposure estimation for the global burden of disease 2013. *Environ Sci Technol* 2016;50(1):79-88. DOI: 10.1021/acs.est.5b03709
73. Anenberg SC, Belova A, Brandt J, Fann N, Greco S, Guttikunda S, Heroux ME, Hurley F, Krzyzanowski M, Medina S, Miller B, Pandey K, Roos J, Van Dingenen R. Survey of ambient air pollution health risk assessment tools. *Risk Anal* 2016 Jan 6. DOI: 10.1111/risa.12540
74. Lentz TJ, Dotson GS, Williams PR, Maier A, Gadagbui B, Pandalai SP, Lamba A, Hearl F, Mumtaz M. Aggregate exposure and cumulative risk assessment-integrating occupational and non-occupational risk factors. *J Occup Environ Hyg* 2015;12(Suppl 1):S112-26. DOI: 10.1080/15459624.2015.1060326
75. Reis S, Seto E, Northcross A, Quinn NW, Convertino M, Jones RL, Maier HR, Schlink U, Steinle S, Vieno M, Wimberly MC. Integrating modelling and smart sensors for environmental and human health. *Environ Model Softw* 2015;74:238-46. DOI: 10.1016/j.envsoft.2015.06.003
76. Cooklin A, Joss N, Husser E, Oldenburg B. Integrated approaches to occupational health and safety: a systematic review. *Am J Health Promot* 2016 DOI: <http://dx.doi.org/10.4278/ajhp.141027-LIT-542>
77. Indinnimeo L, Porta D, Forastiere F, De Vittori V, De Castro G, Zicari A, Tancredi G, Melengu T, Duse M. Prevalence and risk factors for atopic disease in a population of preschool children in Rome: Challenges to early intervention. *Int J Immunopathol Pharmacol* 2016;29(2):308-19. DOI: 10.1177/0394632016635656
78. Boisquet J, Anto JM, Akdis M, Auffray C, Keil T, Momas I, Postma D, Valenta R, Wickman M, Cambon-Thomsen A, Haahtela T, Lambrecht BN, Lodrup-Carlsen K, Koppelman GH, Sunyer J, Zuberbier T, Annesi-Maesano I, Arno A, Bindslev-Jensen C, De Carlo G, Forastiere F, Heinrich J, Kowalski ML, Maier D, Melén E, Palkonen S, Smit HA, Standl M, Wright J, Arsanoj A, Benet M, Balardini N, Garcia-Aymerich J, Gehring U, Guerra S, Hohman C, Kull I, Lupinek C, Pinart M, Skrindo I, Westman M, Smagghe D, Akdis C, Albang R, Anastasova V, Anderson N, Bachert C, Ballereau S, Ballester F, Basagana X, Bedbrook A, Bergstrom A, von Berg A, Brunekreef B, Burte E, Carlsen KH, Chatzi L, Coquet JM, Curin M, Demoly P, Eller E, Fantini MP, Gerhard B, Hammad H, von Hertzen L, Hovland V, Jacquemin B, Just J, Keller T, Kerkhof M, Kiss R, Kogevinas M, Koltzko S, Lau S, Lehmann I, Lemonnier N, McEachan R, Mäkelä M, Mestres J, Minina E, Mowinkel P, Nadif R, Nawijn M, Oddie S, Pellet J, Pin I, Porta D, Rancière F, Rial-Sebbag A, Saes Y, Schuijs MJ, Siroux V, Tischer CG, Torrent M, Varraso R, De Vocht J, Wenger K, Wieser S, Xu C. Paving the way of systems biology and precision medicine in allergic diseases: The MeDALL success story. *Allergy* 2016 Mar 10. DOI: 10.1111/all.12880